

| HEDIS® HYBRID MEASURES | | |
|---|---|--|
| <i>(Hybrid measures are based on data retrieved from medical records and may include administrative data from claims)</i> | | |
| HEDIS® MEASURE | MEMBER DESCRIPTION | HEDIS® MEDICAL RECORD DOCUMENTATION GUIDELINES |
| Adolescent Well-Care Visits (AWC) | Members 12-21 years of age who had either: <ul style="list-style-type: none"> • At least one comprehensive well-care visit with a PCP; or • At least one comprehensive well-care visit with an OB/GYN practitioner | <u>Medical Record must include the following:</u> <ul style="list-style-type: none"> • Date the well-care visit occurred • A health history • A physical developmental history • A mental developmental history • A physical exam • Health education/anticipatory guidance |
| Adult BMI Assessment (ABA) | Members 18-74 years of age who had an outpatient visit and whose: <ul style="list-style-type: none"> • Body Mass Index (BMI) was documented | <u>Medical Record must include the following:</u> <ul style="list-style-type: none"> • Weight and BMI value (for members 20 years of age and older) • Height, weight, and BMI percentile (for members under than 20 years of age); BMI percentile can either be documented as a value or plotted on an age-growth chart |

| HEDIS® MEASURE | MEMBER DESCRIPTION | HEDIS® MEDICAL RECORD DOCUMENTATION GUIDELINES |
|---|--|--|
| <p>Cervical Cancer Screening (CCS)</p> | <p>Women 21-64 years of age who were screened for cervical cancer using either one of the following criteria:</p> <ul style="list-style-type: none"> • <i>Ages 21-64:</i> A cervical cytology every 3 years • <i>Ages 30-64:</i> A cervical cytology/HPV co-testing every 5 years | <p><u>Medical Record must include the following:</u></p> <ul style="list-style-type: none"> • Date the cervical cytology and/or HPV test was performed • The result or finding |
| <p>Childhood Immunization Status (CIS)</p> | <p>Members 2 years of age who had the following vaccines by their 2nd birthday:</p> <ul style="list-style-type: none"> • 4 DTaP • 3 IPV • 1 MMR • 3 HiB • 3 HepB • 1 VZV • 4 PCV • 1 HepA • 2 or 3 RV • 2 Flu | <p><u>Medical Record must include the following:</u></p> <ul style="list-style-type: none"> • A note indicating the name of the specific antigen and the date of the immunization; or • A certificate of immunization prepared by an authorized healthcare provider or agency including the specific dates and types of immunizations administered |

| HEDIS® MEASURE | MEMBER DESCRIPTION | HEDIS® MEDICAL RECORD DOCUMENTATION GUIDELINES |
|---|---|--|
| <p>Comprehensive Diabetes Care (CDC)</p> | <p>Members 18-75 years of age with diabetes (type 1 or 2) who had each of the following:</p> <ul style="list-style-type: none"> • HbA1c testing: <ul style="list-style-type: none"> ○ HbA1c poor control (>9.0%) ○ HbA1c control (<8.0%) ○ HbA1c control (<7.0%) • Retinal eye exam • Medical attention for nephropathy • Blood Pressure (BP) control (<140/90 mm Hg) | <p><u>Medical Record must include the following:</u></p> <ul style="list-style-type: none"> • A note indicating the date when the HbA1c test was performed with the result • Evidence of one of the following: <ul style="list-style-type: none"> ○ A note or letter indicating an ophthalmoscopic exam was completed by an eye care professional, the date the procedure was performed, and the results ○ A chart or photograph of retinal abnormalities indicating the date when the fundus photography was performed with evidence that an eye care professional reviewed the results ○ Documentation of a negative retinal or dilated exam by an eye care professional • Evidence of one of the following: <ul style="list-style-type: none"> ○ A urine test for albumin or protein, the date the test was performed and the result ○ Documentation of a visit to a nephrologist ○ Documentation of a renal transplant ○ Documentation for medical attention for diabetic nephropathy, ESRD, CRF, CKD, renal insufficiency, Proteinuria, Albuminuria, renal dysfunction, ARF, dialysis ○ Evidence of ACE inhibitor/ARB • BP reading |

| HEDIS® MEASURE | MEMBER DESCRIPTION | HEDIS® MEDICAL RECORD DOCUMENTATION GUIDELINES |
|--|---|---|
| <p>Controlling High Blood Pressure (CBP)</p> | <p>Members 18-85 years of age who had a diagnosis of hypertension (HTN) and whose BP was adequately controlled based on the following criteria:</p> <ul style="list-style-type: none"> • <i>Ages 18-59:</i> BP was <140/90 mm Hg • <i>Ages 60-85 with a diagnosis of diabetes:</i> BP was <140/90 mm Hg • <i>Ages 60-85 without a diagnosis of diabetes:</i> BP was <150/90 mm Hg | <p><u>Medical Record must include the following:</u></p> <ul style="list-style-type: none"> • Diagnosis of HTN • Date and result of last BP reading in the measurement year (systolic and diastolic); if multiple readings are present, the lowest systolic and diastolic reading on that date is the representative BP |
| <p>Frequency of Ongoing Prenatal Care (FPC)</p> | <p>Women who delivered a live-birth that had the expected number of prenatal visits based on the month of a member’s enrollment in the HUSKY Health program and gestational age. The following percentages are calculated for this measure:</p> <ul style="list-style-type: none"> • <21% • 21%-40% • 41%-60% • 61%-80% • ≥81% | <p><u>Medical Record must include the following:</u></p> <ul style="list-style-type: none"> • A note indicating the date the prenatal visit occurred • Evidence of one of the following: <ul style="list-style-type: none"> ○ A physical obstetrical exam that includes auscultation for fetal heart tone ○ Pelvic exam with obstetric observations ○ Measurement of fundus height ○ Evidence that a prenatal care procedure was performed (screening test in the form of an obstetric panel, TORCH antibody panel alone, rubella antibody test/titer with an RH incompatibility blood typing, or echography of pregnant uterus) ○ Documentation of LMP or EDD in conjunction with either prenatal risk assessment or complete obstetrical history |

| HEDIS® MEASURE | MEMBER DESCRIPTION | HEDIS® MEDICAL RECORD DOCUMENTATION GUIDELINES |
|--|--|--|
| Immunizations for Adolescents (IMA) | Members 13 years of age who had all of the following by their 13 th birthday: <ul style="list-style-type: none"> • 1 dose of meningococcal vaccine • 1 dose of Tdap • 3 doses of HPV vaccine | <u>Medical Record must include the following:</u> <ul style="list-style-type: none"> • A note indicating the name of the specific antigen and the date of service; or • A certificate of immunization prepared by an authorized healthcare provider or agency including the specific dates and types of immunizations administered |
| Lead Screening in Children (LSC) | Members 2 years of age who had the following by their 2 nd birthday: <ul style="list-style-type: none"> • 1 or more capillary or venous lead blood test for lead poisoning | <u>Medical Record must include the following:</u> <ul style="list-style-type: none"> • A note indicating the date the test was performed • The result or finding |

| HEDIS® MEASURE | MEMBER DESCRIPTION | HEDIS® MEDICAL RECORD DOCUMENTATION GUIDELINES |
|--|---|---|
| <p>Prenatal and Postpartum Care (PPC)</p> | <p>Women who delivered a live-birth that met the following criteria:</p> <ul style="list-style-type: none"> • <i>Timeliness of Prenatal Care:</i> Received a prenatal care visit in the first trimester or within 42 days of enrolling in the HUSKY Health program • <i>Postpartum Care:</i> Had a postpartum visit on or between 21 and 56 days after delivery | <p><u>Medical Record must include the following:</u></p> <p><i>Prenatal Care:</i></p> <ul style="list-style-type: none"> • A note indicating the date the prenatal visit occurred • Evidence of one of the following: <ul style="list-style-type: none"> ○ A physical obstetrical exam that includes auscultation for fetal heart tone ○ Pelvic exam with obstetric observations ○ Measurement of fundus height ○ Evidence that a prenatal care procedure was performed (screening test in the form of an obstetric panel, TORCH antibody panel alone, rubella antibody test/titer with an RH incompatibility blood typing, or echography of pregnant uterus) ○ Documentation of LMP or EDD in conjunction with either prenatal risk assessment or complete obstetrical history <p><i>Postpartum Care:</i></p> <ul style="list-style-type: none"> • A note indicating the date the postpartum visit occurred • Evidence of one of the following: pelvic exam, evaluation of weight, BP, breasts and abdomen, or notation of postpartum care (PP care, PP check, 6-week check, or pre-printed postpartum care form) |

| HEDIS® MEASURE | MEMBER DESCRIPTION | HEDIS® MEDICAL RECORD DOCUMENTATION GUIDELINES |
|---|---|---|
| <p>Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents (WCC)</p> | <p>Members 3-17 years of age who had an outpatient visit with a PCP or OB/GYN and had evidence of the following:</p> <ul style="list-style-type: none"> • BMI percentile documentation • Counseling for nutrition • Counseling for physical activity | <p><u>Medical Record must include the following:</u></p> <ul style="list-style-type: none"> • BMI Percentile: Height, weight and BMI percentile or BMI percentile plotted on an age-growth chart • Counseling for Nutrition*: A note indicating the date counseling occurred and at least one of the following: discussion of current nutrition behaviors, checklist indicating nutrition was addressed, counseling or referral for nutrition education, educational materials on nutrition were provided during a face-to-face visit, anticipatory guidance for nutrition, or weight/obesity counseling • Counseling for Physical Activity*: A note indicating the date counseling occurred and at least one of the following: discussion of current physical activity behaviors, checklist indicating physical activity was addressed, counseling or referral for physical activity, educational materials on physical activity provided during a face-to-face visit, anticipatory guidance specific to the child’s physical activity, or weight/obesity counseling <p><i>*Documentation of obesity counts for both the nutrition and physical activity components of this measure</i></p> |

| HEDIS® MEASURE | MEMBER DESCRIPTION | HEDIS® MEDICAL RECORD DOCUMENTATION GUIDELINES |
|---|---|--|
| <p>Well-Child Visits in the First 15 Months of Life (W15)</p> | <p>Members 15 months old who had the following number of well-child visits with a PCP during their first 15 months of life:</p> <ul style="list-style-type: none"> • 0 visits • 1 visit • 2 visits • 3 visits • 4 visits • 5 visits • 6 or more visits | <p><u>Medical Record must include the following:</u></p> <ul style="list-style-type: none"> • A note indicating a visit with a PCP • The date the well-child visit occurred • Health history • Physical developmental history • Mental developmental history • Complete physical exam • Health education/anticipatory guidance |
| <p>Well-Child Visits in the Third, Fourth, Fifth and Sixth Years of Life (W34)</p> | <p>Members 3-6 years of age who had:</p> <ul style="list-style-type: none"> • 1 or more well-child visits with a PCP | <p><u>Medical Record must include the following:</u></p> <ul style="list-style-type: none"> • A note indicating a visit with a PCP • The date the well-child visit occurred • Health history • Physical developmental history • Mental developmental history • Complete physical exam • Health education/anticipatory guidance |

| HEDIS® ADMINISTRATIVE MEASURES <i>(Administrative measures are based on claims/encounter data)</i> | | |
|--|---|--|
| HEDIS® MEASURE | MEMBER DESCRIPTION | HEDIS® CLAIMS CODING GUIDELINES |
| Adherence to Antipsychotic Medications for Individuals with Schizophrenia (SAA) | Members 19-64 years of age with schizophrenia | Members must have had a diagnosis of schizophrenia and had been: <ul style="list-style-type: none"> • Dispensed and remained on an antipsychotic medication for at least 80% of their treatment period |
| Adults' Access to Preventive/Ambulatory Health Services (AAP) | Members 20 years of age and older | Members must have had: <ul style="list-style-type: none"> • One or more ambulatory or preventive care visit during the measurement year |
| Annual Dental Visit (ADV) | Members 2-20 years of age | Members must have had: <ul style="list-style-type: none"> • At least 1 dental visit during the measurement year |
| Annual Monitoring for Patients on Persistent Medications (MPM) | Members 18 years of age and older who received at least 180 days of ambulatory medication therapy for a select therapy agent, and at least one therapeutic monitoring event for a therapeutic agent | Members must have received 180 treatment days and at least 1 monitoring event for any of the following therapeutic agents: <ul style="list-style-type: none"> • ACE Inhibitors • ARB • Digoxin • Diuretics |

| HEDIS® MEASURE | MEMBER DESCRIPTION | HEDIS® CLAIMS CODING GUIDELINES |
|--|---|---|
| Antidepressant Medication Management (AMM) | Members 18 years of age and older who were treated with an antidepressant and had a diagnosis of major depression | Members must have been treated with an antidepressant, had a diagnosis of major depression, and remained on the antidepressant for the following timeframes: <ul style="list-style-type: none"> • <i>Effective Acute Phase:</i> At least 84 days (3 months) • <i>Effective Continuation Phase:</i> At least 180 days (6 months) |
| Appropriate Testing for Children with Pharyngitis (CWP) | Members 3-18 years of age diagnosed with pharyngitis | Members must have been diagnosed with pharyngitis and have been both: <ul style="list-style-type: none"> • Dispensed an antibiotic • Received a group A strep test |
| Appropriate Treatment for Children with Upper Respiratory Infection (URI) | Members 3 months-18 years of age who were diagnosed with an upper respiratory infection (URI) | Members must have been diagnosed with a URI and: <ul style="list-style-type: none"> • <u>Not</u> dispensed an antibiotic prescription |
| Asthma Medication Ratio (AMR) | Members 5-64 years of age who were diagnosed with persistent asthma | Members must have been diagnosed with persistent asthma and: <ul style="list-style-type: none"> • Had a ratio of controller medications to total asthma medications of 0.50 or greater |
| Avoidance of Antibiotic Treatment in Adults With Acute Bronchitis (AAB) | Members 18-64 years of age who were diagnosed with acute bronchitis | Members must have been diagnosed with acute bronchitis and: <ul style="list-style-type: none"> • <u>Not</u> dispensed an antibiotic treatment |
| Breast Cancer Screening (BCS) | Women 50-74 years of age | Women must have had: <ul style="list-style-type: none"> • One or more mammograms to screen for breast cancer <p><i>Diagnostic mammograms do not meet the guidelines</i></p> |

| HEDIS® MEASURE | MEMBER DESCRIPTION | HEDIS® CLAIMS CODING GUIDELINES |
|---|--|---|
| Cardiovascular Monitoring for People with Cardiovascular Disease and Schizophrenia (SMC) | Members 18-64 years of age with schizophrenia and cardiovascular disease | Members must have had a diagnosis of schizophrenia and cardiovascular disease and had: <ul style="list-style-type: none"> • An LDL-C test |
| Children and Adolescents' Access to Primary Care Practitioners (CAP) | Members 12 months-19 years of age | Members must have had: <ul style="list-style-type: none"> • One or more ambulatory or preventive care visits with a PCP |
| Chlamydia Screening in Women (CHL) | Women 16-24 years of age | Women must have been identified as sexually active and had: <ul style="list-style-type: none"> • At least one test for chlamydia |
| Diabetes Monitoring for People With Diabetes and Schizophrenia (SMD) | Members 18-64 years of age with schizophrenia and diabetes | Members must have been diagnosed with schizophrenia and diabetes and had both: <ul style="list-style-type: none"> • An LDL-C test • An HbA1c test |
| Diabetes Screening for People with Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications (SSD) | Members 18-64 years of age with schizophrenia or bipolar disorder | Members must have been diagnosed with schizophrenia or bipolar disorder and have been both: <ul style="list-style-type: none"> • Dispensed antipsychotic medications • Received a diabetes screening test |

| HEDIS® MEASURE | MEMBER DESCRIPTION | HEDIS® CLAIMS CODING GUIDELINES |
|---|---|---|
| Disease-Modifying Anti-Rheumatic Drug Therapy for Rheumatoid Arthritis (ART) | Members 18 years of age and older who were diagnosed with rheumatoid arthritis | Members must have been diagnosed with rheumatoid arthritis and: <ul style="list-style-type: none"> • Dispensed at least one ambulatory prescription for a disease-modifying anti-rheumatic drug (DMARD) |
| Follow-Up After Emergency Department Visit for Alcohol and Other Drug Dependence (FUA) | The percentage of ED visits for members 13 years of age and older with a principal diagnosis of alcohol or other drug (AOD) dependence who had a follow-up visit for AOD | Members who had an ED visit for alcohol or other drug dependence who had follow-up visits within either: <ul style="list-style-type: none"> • 7 days of the ED visit • 30 days of the ED visit |
| Follow-Up After Emergency Department Visit for Mental Illness (FUM) | The percentage of ED visits for members 6 years of age and older with a primary diagnosis of mental illness, who had an outpatient visit, an intensive outpatient encounter or a partial hospitalization for mental illness | Members who had a follow-up visit after an ED visit for mental illness within either: <ul style="list-style-type: none"> • 7 days of the ED visit • 30 days of the ED visit |
| Follow-Up After Hospitalization for Mental Illness (FUH) | Members 6 years of age and older hospitalized for treatment of selected mental illness | Members must have been hospitalized for treatment of a selected mental illness and had an outpatient visit, intensive outpatient encounter, or partial hospitalization with a mental health practitioner within either: <ul style="list-style-type: none"> • 7 days of discharge • 30 days of discharge |

| HEDIS® MEASURE | MEMBER DESCRIPTION | HEDIS® CLAIMS CODING GUIDELINES |
|--|---|--|
| <p>Follow-Up Care for Children Prescribed ADHD Medication (ADD)</p> | <p>Members 6-12 years of age with newly prescribed ADHD medication</p> | <p>Members must have been newly prescribed an ADHD medication and had:</p> <ul style="list-style-type: none"> • <i>Initiation Phase:</i> 1 follow-up visit with a practitioner of prescribing authority within 30 days • <i>Continuation and Maintenance Phase:</i> Remained on the medication for at least 210 days and, in addition to the Initiation Phase visit, had at least 2 follow-up visits with a practitioner within 270 days (9 months) after the Initiation Phase ended |
| <p>Initiation and Engagement of Alcohol and Other Drug Dependence Treatment (IET)</p> | <p>Members 13 years of age and older with a new episode of alcohol or other drug (AOD) dependence</p> | <p>Members must have had a new episode of alcohol or other drug (AOD) dependence diagnosis and received:</p> <ul style="list-style-type: none"> • <i>Initiation of AOD Treatment:</i> An inpatient AOD admission, outpatient visit, intensive outpatient encounter, or partial hospitalization within 14 days • <i>Engagement of AOD Treatment:</i> Two or more additional services with a diagnosis of AOD within 30 days of the initiation visit |
| <p>Medication Management for People with Asthma (MMA)</p> | <p>Members 5-64 years of age identified as having persistent asthma</p> | <p>Members must have been diagnosed with persistent asthma, dispensed an asthma controller medication and remained on treatment either:</p> <ul style="list-style-type: none"> • At least 50% of their treatment period • At least 75% of their treatment period |

| HEDIS® MEASURE | MEMBER DESCRIPTION | HEDIS® CLAIMS CODING GUIDELINES |
|--|---|--|
| <p>Metabolic Monitoring for Children and Adolescents on Antipsychotics (APM)</p> | <p>Members 1-17 years of age who had two or more antipsychotic prescriptions</p> | <p>Members must have had two or more antipsychotic prescriptions and had metabolic testing including:</p> <ul style="list-style-type: none"> • At least one test for blood glucose • At least one test for LDL-C or cholesterol |
| <p>Non-Recommended Cervical Cancer Screening in Adolescent Females (NCS)*</p> <p><i>*Lower rate indicates better performance.</i></p> | <p>Women 16-20 years of age who were screened unnecessarily for cervical cancer</p> | <p>Evidence-based guidelines indicate that women under 21 years of age should not be screened for cervical cancer regardless of age at sexual initiation or other behavior-related factors. Women screened for either:</p> <ul style="list-style-type: none"> • Cervical cytology • HPV test |
| <p>Persistence of Beta-Blocker Treatment After a Heart Attack (PBH)</p> | <p>Members 18 years of age and older who were hospitalized with a diagnosis of Acute Myocardial Infarction (AMI) and received persistent beta-blocker treatment</p> | <p>Members who were hospitalized and discharged with a diagnosis of AMI who:</p> <ul style="list-style-type: none"> • Received beta-blocker treatment for six months (180 day course) after discharge |
| <p>Pharmacotherapy Management of COPD Exacerbation (PCE)</p> | <p>Members 40 years of age and older who had an acute inpatient discharge or ED visit for COPD exacerbations</p> | <p>COPD exacerbations for members who had an acute inpatient discharge or ED visit who were dispensed either:</p> <ul style="list-style-type: none"> • A systemic corticosteroid within 14 days of the event • A bronchodilator within 30 days of the event |

| HEDIS® MEASURE | MEMBER DESCRIPTION | HEDIS® CLAIMS CODING GUIDELINES |
|---|--|---|
| Statin Therapy for Patients with Cardiovascular Disease (SPC) | Men 21-75 years of age and women 40-75 years of age identified as having clinical atherosclerotic cardiovascular disease (ASCVD) and received statin therapy | Members who were diagnosed as having clinical ASCVD and met the following criteria: <ul style="list-style-type: none"> • <i>Received Statin Therapy</i>: Dispensed at least one high or moderate-intensity statin medication • <i>Statin Adherence 80%</i>: Remained on a high or moderate-intensity statin medication for at least 80% of the treatment period |
| Statin Therapy for Patients with Diabetes (SPD) | Members 40-75 years of age with diabetes who do not have ASCVD and received statin therapy | Members who were diagnosed with diabetes and did not have ASCVD and met the following criteria: <ul style="list-style-type: none"> • <i>Received Statin Therapy</i>: Dispensed at least one statin medication of any intensity • <i>Statin Adherence 80%</i>: Remained on a statin medication of any intensity for at least 80% of the treatment period |
| Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics (APP) | Members 1-17 years of age with a new prescription for an antipsychotic medication | Members who had a new prescription for an antipsychotic medication and had: <ul style="list-style-type: none"> • Documentation of psychosocial care as first-line treatment in the 120-day period from 90 days prior to the Index Prescription Start Date (IPSD) through 30 days after the IPSD |
| Use of Imaging Studies for Low Back Pain (LBP) | Members 18-50 years of age with a primary diagnosis of low back pain | Members who were diagnosed with low back pain and: <ul style="list-style-type: none"> • Did <u>not</u> have an imaging study (X-ray, MRI, CT Scan) within 28 days of the diagnosis |

| HEDIS® MEASURE | MEMBER DESCRIPTION | HEDIS® CLAIMS CODING GUIDELINES |
|--|--|---|
| <p>Use of Multiple Concurrent Antipsychotics in Children and Adolescents (APC)*</p> <p><i>*Lower rate indicates better performance.</i></p> | <p>Members 1-17 years of age</p> | <p>Members who were on:</p> <ul style="list-style-type: none"> • Two or more concurrent antipsychotic medications |
| <p>Use of Spirometry Testing in the Assessment and Diagnosis of COPD (SPR)</p> | <p>Members 40 years of age and older with a new diagnosis of COPD or newly active COPD</p> | <p>Members who were newly diagnosed with COPD and who received:</p> <ul style="list-style-type: none"> • Spirometry testing to confirm the diagnosis |